

$\mathbb{Q}$ -factorial quartic threefolds

C. SHRAMOV

ABSTRACT. We prove that a nodal quartic threefold X containing no planes is $\mathbb{Q}$ -factorial provided that it has not more than 12 singular points, with the exception of a quartic with exactly 12 singularities containing a quadric surface. We give some geometrical constructions related to the latter quartic.

1. INTRODUCTION

A quartic threefold is a classical object of interest in birational geometry. The group of birational automorphisms of a nonsingular quartic was first studied in [2]; birational automorphism groups of singular quartics were studied in particular in [4] (the case of a single simple double point) and [14] (the case of arbitrary number of simple double points). M. Mella (see [14]) proved birational rigidity and described the generators of the group of birational automorphisms of any $\mathbb{Q}$ -factorial nodal quartic. On the other hand, one cannot extend these results to the case of non- $\mathbb{Q}$ -factorial varieties: there are non- $\mathbb{Q}$ -factorial nodal quartics threefolds that are rational. This motivates the following question: which conditions guarantee $\mathbb{Q}$ -factoriality of a nodal quartic?

Recall that a variety V is called *factorial* if any Weil divisor on V is also a Cartier divisor, and *$\mathbb{Q}$ -factorial* if an appropriate multiple of any Weil divisor is a Cartier divisor. A singular point $x \in V$ is called a *simple double point* (or a *simple double singularity*) if it has a neighborhood analytically isomorphic to a neighborhood of a vertex of a cone over a nonsingular quadric. If V is a hypersurface in $\mathbb{P}^n$ given by the equation $F = 0$, the latter is equivalent to non-degeneracy of the Hessian $H(F)$ at x . The variety with only simple double singularities is called *nodal*.

Let X be a nodal quartic threefold. Then the following is well-known.

Theorem 1.1 (see [7]). *If the number of singular points of X does not exceed 8, then X is $\mathbb{Q}$ -factorial; if it is equal to 9, and X contains no planes, then X is again $\mathbb{Q}$ -factorial.*

This result is a particular case of the following conjecture (see, for example, [10], [7], and also [8] for some advances).

Conjecture 1.2. *Let Y be a nodal hypersurface of degree d in $\mathbb{P}^4$ with singular points. If $s < (d - 1)^2$, then Y is $\mathbb{Q}$ -factorial; if Y contains no planes and $s < 2(d - 1)(d - 2)$, then Y is $\mathbb{Q}$ -factorial; if $s \leq 2(d - 1)(d - 2)$, and Y contains neither planes nor quadrics, then Y is again $\mathbb{Q}$ -factorial.*

By now Conjecture 1.2 is checked only for $d \leq 3$.

The main goal of this paper is to show the following.

Theorem 1.3. *Let a nodal quartic threefold X with not more than 11 singularities contain no planes. Then X is $\mathbb{Q}$ -factorial. If X has 12 singularities, then X is again $\mathbb{Q}$ -factorial with the exception of the case when X contains a quadric surface.*

Theorem 1.3 generalizes Theorem 1.1; it is a particular case of Conjecture 1.2. Since there are non- $\mathbb{Q}$ -factorial nodal quartics containing no planes and having 12 singular points (the example is given by a general quartic containing a quadric — see Example 4.1 below), the statement of Theorem 1.3 cannot be refined (as well as the general statement of Conjecture 1.2).

The proof of Theorem 1.3 occupies section 3; it uses some simple consequences of Bese theorem (see Theorem 2.1) that are collected in section 2. In section 4 we consider some constructions related to a general quartic containing a quadric: we describe a (well-known) transformation of this quartic to a complete intersection of a quadric and a cubic in $\mathbb{P}^5$ with a single double point (Example 4.1); we also give an example of a birational involution of the latter variety that does not exist in the nonsingular case (Example 4.4 and Statement 4.5).

All the varieties are defined over the field of complex numbers $\mathbb{C}$.

The author is grateful to I. Cheltsov and S. Galkin for useful discussions.

2. AUXILIARY STATEMENTS ABOUT POINTS ON SURFACES

In this section we collect some statements that will be used in section 3.

Theorem 2.1 (see [5]). *Let $\pi : \mathbb{F}_{r,N} \rightarrow \mathbb{F}_r$ be a blow-up of a rational ruled surface $\mathbb{F}_r$ at N points $p_1, \dots, p_N$. Let $D \sim as + bf$ be a divisor on $\mathbb{F}_r$, where s and f denote the classes of an exceptional section and a fiber of $\mathbb{F}_r$ respectively, and $a, b > 0$. Let $L_N = \pi^*D - \sum_{i=1}^N E_i$, where E_i are exceptional divisors of π . Let $\rho = \dim |D - K_{\mathbb{F}_r}|$, and*

$$h = h^1(\mathbb{F}_r, \mathcal{O}_{\mathbb{F}_r}(D) \otimes K_{\mathbb{F}_r}^{-1}) = \rho - \frac{D^2 - 3DK_{\mathbb{F}_r} + 16}{2}.$$

Then the line bundle $\mathcal{O}_{\mathbb{F}_r, N}(L_N)$ is generated by its global sections, provided that the following conditions hold:

- (i) $N \leq \lfloor \frac{\rho-4}{3} \rfloor$,
- (ii) $D^2 \geq 7 + 4h$,
- (iii) any curve C of bidegree $(x, y) \neq (0, 0)$ (i. e. linearly equivalent to $xs + yf$), $0 \leq x \leq a + 2$, $0 \leq y \leq \lfloor \frac{b+2+r}{2} \rfloor$, contains not more than $C(D - K_{\mathbb{F}_r} - C) - 2$ points of the set $\{p_1, \dots, p_N\}$.

Corollary 2.2. *Let $p_1, \dots, p_{10}, q \in \mathbb{F}_0 = \mathbb{P}^1 \times \mathbb{P}^1$ be such points that no curve of bidegree $(1, 0)$ or $(0, 1)$ contains 4 points of $p_1, \dots, p_{10}$, no curve of bidegree $(1, 1)$ contains 7 of them and no curve of bidegree $(2, 1)$ or $(1, 2)$ contains all of them. Then there is a divisor D of bidegree $(3, 3)$ passing through $p_1, \dots, p_{10}$ and not passing through q .*

Proof. In the notations of Theorem 2.1 we have $\rho = 35$, $h = 0$, $D^2 = 18$, hence the conditions (i) and (ii) hold. Condition (iii) claims that any curve C of bidegree $(x, y) \neq (0, 0)$, $0 \leq x \leq 5$, $0 \leq y \leq 2$, contains not more than

$$C(D - K_{\mathbb{P}^1 \times \mathbb{P}^1} - C) - 2 = 5x + 5y - 2xy - 2$$

points of $p_1, \dots, p_{10}$; the check of the latter is straightforward for all the pairs (x, y) . $\square$

Corollary 2.3. *Let $p_1, \dots, p_{10}, q \in \mathbb{F}_2$ be such points that the exceptional section contains no points of $p_1, \dots, p_{10}$, no fiber contains 4 of them, no curve of bidegree $(1, 2)$ contains 7 of them and no curve of bidegree $(1, 3)$ contains all of them. Then there is a divisor D of bidegree $(3, 6)$ passing through $p_1, \dots, p_{10}$ and not passing through q .*

Proof. In the notations of Theorem 2.1 we have $\rho = 35$, $h = 0$, $D^2 = 18$, hence the conditions (i) and (ii) hold. Condition (iii) claims that any curve C of bidegree $(x, y) \neq (0, 0)$, $0 \leq x \leq 5$, $0 \leq y \leq 5$, contains not more than $2x^2 + 5y - 2xy - 2$ points of $p_1, \dots, p_{10}$ and is a straightforward consequence of the assumptions (note that condition (iii) for $(x, y) = (1, 0)$ means that the exceptional section does not contain any of the points p_i). $\square$

Corollary 2.4. *Let $p_1, \dots, p_8, q \in \mathbb{F}_2$ be such points that the exceptional section contains no points of $p_1, \dots, p_8$, no fiber contains 4 of them, no curve of bidegree $(1, 2)$ contains 6 of them and no curve of bidegree $(1, 3)$ contains all of them. Then there is a divisor D of bidegree $(2, 5)$ passing through $p_1, \dots, p_8$ and not passing through q .*

Proof. In the notations of Theorem 2.1 we have $\rho = 29$, $h = 0$, $D^2 = 12$, hence conditions (i) and (ii) hold. Condition (iii) claims that any curve

C of bidegree $(x, y) \neq (0, 0)$, $0 \leq x \leq 4$, $0 \leq y \leq 4$, contains not more than $2x^2 + x + 4y - 2xy - 2$ points of $p_1, \dots, p_8$ and is a straightforward consequence of the assumptions. $\square$

Corollary 2.5. *Let $Y \subset \mathbb{P}^3$ be an irreducible quadric; let $p_1, \dots, p_{10}, q \in Y$ be such points that no line contains 4 of the points $p_1, \dots, p_{10}$ no conic contains 7 of them and no twisted cubic contains all of them. Then there is a divisor $D \in \mathcal{O}_{\mathbb{P}^3}(3)|_Y$ passing through $p_1, \dots, p_{10}$ and not passing through q .*

Proof. If Y is a nonsingular quadric, then the statement follows from Corollary 2.2. If Y is a cone over a conic and its vertex does not coincide with any of the points p_i, q , then we are done by Corollary 2.3 applied to the preimages of p_i, q under the resolution map $g : \mathbb{F}_2 \rightarrow Y$. If q is the vertex, we apply Corollary 2.3 to the points $g^{-1}(p_i)$ and an arbitrary point $\tilde{q} \in s \subset \mathbb{F}_2$; this gives us a divisor $\tilde{D} \sim 3s + 6f$ passing through $g^{-1}(p_i)$ and not containing the exceptional section s , and hence not intersecting s , since $\tilde{D}s = 0$. The divisor $D = g_*\tilde{D}$ is just what we looked for.

Now we have to consider the case when one of the points p_i (say, p_{10}) coincides with the vertex of Y . Since not all of the points p_i lie on a single line, we may choose a line $l \subset Y$ such that $q \notin l$ and one of the points $p_1, \dots, p_9$ is contained in l . Moreover, if there is a conic containing 6 of the points $p_1, \dots, p_9$, then such a conic is unique, and we can choose l so that it contains one of these 6 points (say, p_9 again). So we may assume that no 3 of the points of $p_1, \dots, p_8$ lie on a line, no 6 of them lie on a conic and all 8 are not contained in a twisted cubic. Applying Corollary 2.4 to the points $g^{-1}(p_1), \dots, g^{-1}(p_8), g^{-1}(q)$, we find a divisor $\tilde{D} \sim 2s + 5f$ passing through $g^{-1}(p_1), \dots, g^{-1}(p_8)$ and not passing through $g^{-1}(q)$. Now we construct the desired divisor as $D = g_*\tilde{D} \cup l$. $\square$

3. $\mathbb{Q}$ -FACTORIALITY

Throughout this section X denotes a nodal quartic threefold containing no plains and singular at the points $p_1, \dots, p_s$.

In the case of a nodal threefold $\mathbb{Q}$ -factoriality is equivalent to factoriality; on the other hand, factoriality of a nodal Fano threefold X is equivalent to a topological condition $\text{rk}H^2(X, \mathbb{Z}) = \text{rk}H_4(X, \mathbb{Z})$ (in our case $\text{rk}H^2(X, \mathbb{Z}) = \rho(X) = 1$). Hence to prove $\mathbb{Q}$ -factoriality of X it suffices to check that for a small resolution $h : \tilde{X} \rightarrow X$ we have $\rho(\tilde{X}) = s + 1$. To check this it is sufficient to show that, using the terminology of [11], the defect of X equals zero (cf. the proof of Theorem 2

in [11], and also [9]). The latter condition means that the singular points $p_1, \dots, p_s$ impose independent conditions on the hypersurfaces of degree 3 in $\mathbb{P}^4$, i.e. that for any point p_i , $1 \leq i \leq s$, there is a cubic hypersurface $D_i \subset \mathbb{P}^4$ such that $p_i \notin D_i$ and $p_j \in D_i$ for $j \neq i$.

The main tool for checking this independency condition is provided by the following theorem proved in [12].

Theorem 3.1. *The points $p_1, \dots, p_s \in \mathbb{P}^n$ impose independent conditions on forms of degree d if each linear subspace of dimension k contains at most $dk + 1$ of the points $p_1, \dots, p_s$.*

The points $p_1, \dots, p_s$ have the following property.

Lemma 3.2. *No line contains 4 of the points $p_1, \dots, p_s$, no plane contains 7, and no twisted cubic contains 10 of them. If the points $p_1, \dots, p_6$ are contained in a plane P , then either X intersects P along a (possibly reducible) double conic, or X intersects P along a union of four different lines $l_1, \dots, l_4$ and the points p_i , $1 \leq i \leq 6$, coincide with the points of pairwise intersections of the lines l_j, l_k .*

Proof. The first three statements hold regardless to the assumption that X contains no planes (see, for example, [8, Lemma 2.7]). Let the points $p_1, \dots, p_6$ lie on a plane P . Assume that $p_1, \dots, p_6$ lie on a conic Q . If there exists a point $q \in (X \cap P) \setminus Q$, then there is a two-dimensional family of plane cubics passing through $p_1, \dots, p_6, q$. Let C be a general curve of this family. Then C is irreducible and $C \subset X$, since otherwise the intersection index of C and X in $\mathbb{P}^4$ would be at least 13. As C is movable, we have $P \subset X$ that is a contradiction.

Now assume that $X \cap P$ is neither a conic nor a union of four lines. It means that there is a point $q \in X \cap P$ such that the points $p_1, \dots, p_6, q$ are not contained in a single conic, and no 4 of them are collinear. So there is again a two-dimensional family of plane cubics passing through $p_1, \dots, p_6, q$; a general curve of this family is irreducible and hence is contained in X , a contradiction. $\square$

Theorem 3.1 and Lemma 3.2 immediately imply

Corollary 3.3. *Let $s \leq 10$. Then X is $\mathbb{Q}$ -factorial.*

On the other hand, Lemma 3.2 gives the following

Corollary 3.4. *Assume that $s \geq 11$ and the points $p_1, \dots, p_s$ are contained in a reducible quadric surface (i.e. in a pair of planes which span a three-dimensional subspace). Then either $p_1, \dots, p_s$ impose independent conditions on the forms of degree 3, or $p_1, \dots, p_s$ are also contained in an irreducible quadric surface.*

Proof. Let $p_1, \dots, p_s$ be contained in a union of the planes L and L' so that $p_1, \dots, p_6 \in L$, $p_7, \dots, p_s \in L'$, $L \cap L' = l$. Note that $s \leq 12$, and at most one of the points p_i lie on the line l . In particular, $p_1, \dots, p_6$ are contained in a conic if and only if $p_7, \dots, p_s$ are, since our assumptions imply that the union of four lines that intersect in singular points of X intersects the line l at least by three points, and a conic intersects l at most by two points.

Assume that the points $p_1, \dots, p_6$ lie on a conic Q_1 , and the points $p_7, \dots, p_s$ lie on a conic Q_2 . Note that $Q_1 \cap l = Q_2 \cap l$ (otherwise X would contain one of the planes L or L'), i.e. the conics Q_i intersect l by the same pair of (possibly coinciding) points. Any two conics with such a property are contained in an irreducible quadric.

Hence if the points $p_1, \dots, p_s$ are not contained in an irreducible quadric, Lemma 3.2 implies that the intersection of X and L is a union of four lines $l_1, \dots, l_4$. So the intersection of X and L' is also a union of four lines; denote them by $l'_1, \dots, l'_4$. We may assume (renumbering the lines l'_i if necessary) that $l_i \cap l = l'_i \cap l$. Now it is easy to see that the points $p_1, \dots, p_s$ impose independent conditions on the forms of degree 3. (For example, to construct a cubic surface S_1 passing through $p_2, \dots, p_s$ and not passing through p_1 , we may proceed as follows: if p_1 is not contained in l_1 and l_2 , then S_1 is a union of the planes $P_1 = \langle l_1, l'_1 \rangle$, $P_2 = \langle l_2, l'_2 \rangle$ and some plane P_3 , passing through the only point of $p_7, \dots, p_s$ that lies on neither l'_1 nor l'_2 , and passing through no other singular point.) $\square$

We'll also need the following observations.

Lemma 3.5. *Assume that $s \leq 12$ and the points $p_1, \dots, p_s$ are contained in a three-dimensional subspace $\mathbb{P}^3 \subset \mathbb{P}^4$ but not in a quadric surface. Then $p_1, \dots, p_s$ impose independent conditions on the forms of degree 3 in $\mathbb{P}^3$ (and hence in $\mathbb{P}^4$ as well).*

Proof. We may assume that $s = 12$. It suffices to find a cubic surface, passing through $p_1, \dots, p_{11}$ and not passing through p_{12} . Since the points $p_1, \dots, p_{12}$ are not contained in a single quadric, we may find a quadric containing 9 of them (say, $p_1, \dots, p_9$) and not containing p_{12} . If the points p_{10} , p_{11} and p_{12} are noncollinear, we may construct the desired cubic surface as a union of this quadric with a plane passing through p_{10} and p_{11} and not passing through p_{12} . If the points p_{10} , p_{11} and p_{12} are collinear, Lemma 3.2 implies that the points p_i , p_{11} and p_{12} are not collinear for any $1 \leq i \leq 9$. On the other hand, not all the quadrics passing through the collections of the points

$\{p_1, \dots, p_9, p_{10}\} \setminus \{p_i\}$, $1 \leq i \leq 9$, also pass through p_{12} , and so we may again obtain our cubic as a union of a quadric and a plane. $\square$

Lemma 3.6. *Assume that $s \leq 11$ and the points $p_1, \dots, p_s$ are contained in a quadric surface. Then $p_1, \dots, p_s$ impose independent conditions on the forms of degree 3 in $\mathbb{P}^4$.*

Proof. Due to Corollary 3.4 we may assume that the points $p_1, \dots, p_s$ are contained in an irreducible quadric. Corollary 2.5 finishes the proof. $\square$

Corollary 3.7. *Assume that $s = 12$ and the points $p_1, \dots, p_{11}$ span a three-dimensional subspace not containing p_{12} . Then $p_1, \dots, p_{12}$ impose independent conditions on the forms of degree 3 in $\mathbb{P}^4$.*

Proof. Apply Lemma 3.5 and Lemma 3.6. $\square$

Lemma 3.8. *Assume that $s = 12$ and the points $p_1, \dots, p_{12}$ are contained in a quadric surface. Then either $p_1, \dots, p_{12}$ impose independent conditions on the forms of degree 3 in $\mathbb{P}^4$, or $p_1, \dots, p_{12}$ are contained in a pencil of quadric surfaces in some three-dimensional subspace.*

Proof. Due to Corollary 3.4 we may assume that the points $p_1, \dots, p_s$ are contained in an irreducible quadric Q ; we may also assume that $s = 12$. Now assume that any quadric passing through $p_1, \dots, p_{12}$ coincides with Q (otherwise the second assertion holds). Let us prove that there exists a (reducible) cubic surface passing through $p_1, \dots, p_{11}$ and not passing through p_{12} . Let Q' be a quadric different from Q , such that Q' contains 8 of the points p_i (say, $p_1, \dots, p_8$) and does not contain p_{12} . If the points $p_9, \dots, p_{12}$ are noncoplanar, we may construct the cubic surface in question as a union of a quadric Q' and a plane passing through p_9, p_{10} and p_{11} . So let the points $p_9, \dots, p_{12}$ lie in a plane P ; renumbering the points p_9, p_{10}, p_{11} if necessary, we may assume that P is spanned by p_{10}, p_{11} and p_{12} . By Lemma 3.2 this plane contains at most two of the points $p_1, \dots, p_8$. Let these be p_7 and p_8 (a similar argument is applicable to the cases when P contains either only one of the points $p_1, \dots, p_8$ or none of them). If all the quadrics passing through the collections $\{p_1, \dots, p_8, p_9\} \setminus \{p_i\}$, $1 \leq i \leq 6$, also pass through the point p_{12} , then the following condition holds: any quadric, containing the points p_7, p_8 and p_9 also contains p_{12} , i.e. the points p_7, p_8, p_9 and p_{12} are collinear; the latter contradicts Lemma 3.2. Otherwise there exists, say, a quadric Q_1 containing $p_2, \dots, p_9$ and not containing p_{12} ; now we obtain a desired cubic as a union of Q_1 with a plane passing through p_1, p_{10} and p_{11} . $\square$

Lemma 3.9. *Assume that $s = 12$ and the points $p_1, \dots, p_{12}$ are contained in a pencil of quadric surfaces in some three-dimensional subspace. Then X contains a quadric surface.*

Proof. Let Q and Q' be two different quadric surfaces containing $p_1, \dots, p_{12}$; let a curve E be the intersection of Q and Q' . We may assume that Q and Q' are irreducible and Q contains a point $q \in X$, $q \notin E$. Let $\mathcal{H}$ be a linear system on Q that consists of the restrictions of the cubic forms vanishing at the points $p_1, \dots, p_{12}$ and q . Lemma 3.2 implies that $\mathcal{H}$ has no base components. But the intersection index of the corresponding curves with X in $\mathbb{P}^4$ is at least 25, hence $Q \subset X$. $\square$

Corollary 3.10. *Assume that $s = 12$ and 11 of the points $p_1, \dots, p_{12}$ are contained in a three-dimensional subspace. Then X is either $\mathbb{Q}$ -factorial, or contains a quadric.*

Proof. Apply Corollary 3.7, Lemma 3.5, Lemma 3.8 and then Lemma 3.9. $\square$

Proof of Theorem 1.3. Due to Corollary 3.3 we may assume that $11 \leq s \leq 12$. Let $s = 11$. If the points $p_1, \dots, p_{11}$ span a three-dimensional subspace, they impose independent conditions on forms of degree 3 in $\mathbb{P}^4$ by Lemma 3.6 and Lemma 3.5. If they span the whole $\mathbb{P}^4$, the statement follows from Theorem 3.1 and Lemma 3.2.

Now let $s = 12$. If no 11 of the points $p_1, \dots, p_{12}$ are contained in a three-dimensional subspace, then $\mathbb{Q}$ -factoriality of X is again implied by Theorem 3.1. If 11 of these points are contained in a three-dimensional subspace, we are done by Corollary 3.10. $\square$

4. SOME CONSTRUCTIONS

Example 4.1 (see, for example, [7, Example 3], [8, Example 1.21], and [14, Example 6]). Let X be a quartic in $\mathbb{P}^4$ with homogeneous coordinates $(x_0 : \dots : x_4)$ containing an irreducible quadric. Then X is described by an equation $QQ' - LC = 0$, where $\deg(Q) = \deg(Q') = 2$, $\deg(L) = 1$, $\deg(C) = 3$ (for brevity we'll also denote the corresponding hypersurfaces in $\mathbb{P}^4$ by the symbols L and C and the corresponding quadric surfaces by the symbols Q and Q'). A sufficiently general quadric of this kind has 12 simple double singularities and is apparently non- $\mathbb{Q}$ -factorial.

Let $y = Q/L = C/Q'$, $y' = Q'/L = C/Q$. A rational map

$$f : (x_0 : \dots : x_4) \mapsto (x_0 : \dots : x_4 : y)$$

gives a birational equivalence of X and a variety $Y \subset \mathbb{P}^5_{(x_0:\dots:x_4:y)}$ given by the equations

$$\begin{aligned} yL &= Q, \\ yQ' &= C. \end{aligned}$$

Similarly a map $f' : (x_0 : \dots : x_4) \mapsto (x_0 : \dots : x_4 : y')$ transforms X to a variety $Y' \subset \mathbb{P}^5_{(x_0:\dots:x_4:y')}$ given by the equations

$$\begin{aligned} y'L &= Q', \\ y'Q &= C. \end{aligned}$$

The map f (resp., f') is a composition of one of the two small resolutions of X , i. e. the blow-up of the (non-Cartier) divisor Q (resp., Q'), and a contraction of the strict transform of the divisor Q' (resp., Q) to a singular point on Y . If Q' is nonsingular, then the corresponding point on Y is a simple double point. A composition $f' \circ f^{-1} : Y \dashrightarrow Y'$ is a link of type II (see [13] for a definition). Varieties Z and Z' are connected by a flop χ .

$$\begin{array}{ccccc} & Z & \overset{\chi}{\dashrightarrow} & Z' & \\ \sigma \swarrow & & & & \searrow \sigma' \\ Y & & & & Y' \\ & \searrow \phi & & \swarrow \phi' & \\ & X & & & \end{array}$$

Similarly, if we have a general intersection Y of a quadric D_2 and a cubic D_3 in $\mathbb{P}^5$ with a unique simple double point p , then Y contains 12 lines $l_1, \dots, l_{12}$ passing through p , all of them contained in the common tangent space $\tilde{L}$ to D_2 and D_3 at p (the generality condition lets us assume that both D_2 and D_3 are smooth). Projection $\pi : Y \dashrightarrow \mathbb{P}^4$ from the point p is a composition of a blow-up $\sigma : Z \rightarrow Y$ of the point p and a contraction $\phi : Z \rightarrow X$ of the preimages of the 12 lines. It is easy to see that the image X of the variety Y under this projection is a quartic, that it contains a quadric $\pi(\tilde{L} \cap D_2)$ and that it is singular exactly at the 12 points $\pi(l_i)$.

Remark 4.2. If the quadric D_2 from Example 4.1 degenerates to a cone over a smooth three-dimensional quadric with a vertex at p , and the cubic D_3 passes through p and remains sufficiently general, then Y has a unique simple double point and is birationally equivalent to a non- $\mathbb{Q}$ -factorial double quadric ramified over a quartic section; the double cover structure is given by the projection π . Note that this gives

an example of a non- $\mathbb{Q}$ -factorial nodal double quadric ramified over a quartic section with the least possible number of singularities.

Remark 4.3 (cf. [6, Conjecture 9]). If the quartic X containing a quadric is sufficiently general, then it has no (biregular) automorphisms interchanging the quadrics Q and Q' . Indeed, each automorphism of X preserves the elliptic curve $E = Q \cap Q'$ and acts on a set $\mathcal{Q} = \{q_1, \dots, q_{16}\}$ of points $q_i \in E$ such that $4q_i$ is a hyperplane section of E ; if the quadrics Q and Q' are sufficiently general, then all the fixed points of the automorphisms from $\text{Aut}(E)$, acting on $\mathcal{Q}$, are contained in $\mathcal{Q}$. Let $\tau_1, \dots, \tau_N \in \text{PGL}_4$ be all automorphisms of $\mathbb{P}^3$ that act on the set $\mathcal{Q}$ and have no fixed points on $E \setminus \mathcal{Q}$; for $t \in E \setminus \mathcal{Q}$ let $t_j = \tau_j(t)$, $1 \leq j \leq N$. Choose a cubic C so that C passes through t but not through any of the points t_j . Assume that there is a nontrivial automorphism $\tau \in \text{Aut}(X)$. Let $\tau' \in \text{PGL}_4$ be such an automorphism of the subspace L that $\tau|_L = \tau'|_L$. Then τ' must coincide with one of the automorphisms τ_j . Moreover, it must act on the set of the singular points of X , and one of these is t . Hence $\tau' = \text{id}$.

Now let X be a quartic without automorphisms interchanging Q and Q' . Then the varieties Y and Y' are not isomorphic since any isomorphisms of Y and Y' gives rise to an automorphism of X interchanging Q and Q' . Hence a general complete intersection of a quadric and a cubic in $\mathbb{P}^5$ with one singular point is not birationally rigid. On the other hand, one can easily construct an example of a quartic X corresponding to isomorphic Y and Y' (see [14, Example 6]).

A complete intersection of a quadric and a cubic in $\mathbb{P}^5$ is a popular object of study. A general variety $\mathcal{V}$ of this type is proved to be birationally rigid, and the group of its birational automorphisms $\text{Bir}(\mathcal{V})$ is completely described (see [1] and [3]). This group is a semidirect product of its subgroup of biregular automorphisms $\text{Aut}(\mathcal{V})$ and a subgroup generated by birational involutions corresponding to lines and conics with linear spans contained in the quadric. The former involutions come from the double cover structures, and the latter correspond to elliptic fibrations. The following example shows that there are some more birational involutions in the singular case.

Example 4.4. Let $B \subset Y$ be a conic passing through the singular point p such that its linear span is not contained in D_2 (a general Y contains 240 conics of this type — see Remark 4.6). A general three-dimensional subspace containing B intersects D_2 by a nonsingular quadric S , and B is a curve of bidegree $(1, 1)$ on S . The intersection of D_3 with S consists of B and an elliptic curve B' of bidegree $(2, 2)$, passing through p and intersecting B by three more points. Let

$g : \tilde{Y} \rightarrow Y$ be a composition of blow-ups of p and of a strict transform of the conic B ; let E and F be the corresponding exceptional divisors. Then $\tilde{Y}$ is endowed by a structure of an elliptic fibration $\phi : \tilde{Y} \rightarrow \mathbb{P}^2$ with a section E and a 3-section F . A reflection with respect to the section E (defined at least on smooth fibers) gives rise to a birational automorphism $\tilde{\tau}_B$ of $\tilde{Y}$ and to a birational involution τ_B of Y . The action of $\tilde{\tau}_B$ on $\text{Pic}(\tilde{Y})$ is described by Statement 4.5.

Statement 4.5 (cf. [3, Lemma 5.1.3]). *Let h , f and e denote the classes of H , F and E in $\text{Pic}(\tilde{X})$. Then*

$$\begin{aligned}\tilde{\tau}_B^* h &= 15h - 8f - 16e, \\ \tilde{\tau}_B^* f &= 14h - 7f - 16e, \\ \tilde{\tau}_B^* e &= e.\end{aligned}$$

Proof. The third equality is obvious. Let G be a general fiber of the fibration ϕ ; let h_G , f_G and e_G denote the restrictions of h , f and e to G . Note that $\tilde{\tau}_B$ acts on $\text{Pic}(G)$, and

$$\tilde{\tau}_B^* h_G = 8f_G - h_G + m(h_G - f_G - e_G)$$

for some $m \in \mathbb{Z}$ since the kernel of the restriction map $\text{Pic}(\tilde{X}) \rightarrow \text{Pic}(G)$ is generated by the class $h_G - f_G - e_G$.

Let $S = \phi^{-1}(l)$ denote a preimage of a general line $l \subset \mathbb{P}^2$; let h_S , f_S and e_S be restrictions of h , f and e to S . Then

$$\begin{aligned}h_S^2 &= 4, h_S f_S = 0, h_S e_S = 2, \\ f_S^2 &= e_S^2 = -2, f_S e_S = 1.\end{aligned}$$

Since $(\tilde{\tau}_B^* h_S)^2 = h_S^2 = 6$, we have

$$6 = ((8f_S - h_S) + m(h_S - f_S - e_S))^2 = -122 + 8m,$$

and hence $m = 16$ that gives the first equality.

Finally, the second inequality is implied by the other two since the involution $\tilde{\tau}_B$ preserves the class $h - e - f$. $\square$

Remark 4.6. If X is sufficiently general, then through any of its singular points p_i , $i = 1, \dots, 12$, there pass 24 lines $l_{i,j}$, $j = 1, \dots, 24$; two of them (say, $l_{i,1}$ and $l_{i,2}$) are contained in the quadric Q' , two (say, $l_{i,3}$ and $l_{i,4}$) are contained in the quadric Q , and other 20 are not contained even in the subspace L . Each of the points p_i corresponds to a birational involution σ_i of X , and each line $l_{i,j}$ corresponds to a birational involution $\sigma_{i,j}$ (see [4] or [3]). On the other hand, each conic $B \subset Y$ such that its linear span is contained in D_2 corresponds to a birational involution τ_B of Y (see. [1] or [3]); each conic $B \subset Y$ such

that its linear span is not contained in D_2 , and $p \in B$, also corresponds to some birational involution τ_B (see Example 4.4). The images of the lines $l_{i,3}$ and $l_{i,4}$ under the map f are conics $B_{i,3}$, $B_{i,4}$ with linear spans contained in D_2 , and the images of the lines $l_{i,j}$, $5 \leq j \leq 24$, are conics $B_{i,j}$ with linear spans not contained in D_2 . All the conics $B_{i,j}$, $1 \leq i \leq 12$, $3 \leq j \leq 24$, pass through the singular point p . Finally, let $l_i \subset Y$ be a line such that $\phi(l_i) = p_i$; each line $l \subset Y$ corresponds to a birational involution τ_l (see [1] or [3]). Under the natural isomorphism of the groups $\text{Bir}(X)$ and $\text{Bir}(Y)$ the involutions σ_i are identified with the involutions τ_{l_i} , $1 \leq i \leq 12$, and the involutions $\sigma_{i,j}$ are identified with the involutions $\tau_{B_{i,j}}$, $1 \leq i \leq 12$, $3 \leq j \leq 24$.

REFERENCES

- [1] V. A. Iskovskikh, *Birational automorphisms of three-dimensional algebraic varieties*, Itogi Nauki Tekh. Sovrem. Probl. Mat., vol. 12, Moscow, VINITI, 1979, 159–235; English transl.: J. Soviet Math., **13** (1980), 815–867.
- [2] V. A. Iskovskikh, Yu. I. Manin, *Three-dimensional quartics and counterexamples to the Lüroth problem*, Mat. Sb., 1971, **86**, 1, 140–166; English transl.: Math. USSR-Sb., 1971, **15**, 1, 141–166.
- [3] V. A. Iskovskikh, A. V. Pukhlikov, *Birational automorphisms of multidimensional algebraic varieties*, Itogi Nauki Tekh. Sovrem. Probl. Mat., vol. 19, Moscow, VINITI, 2001, 5–139.
- [4] A. V. Pukhlikov, *Birational automorphisms of three-dimensional quartic with an elementary singularity*, Mat. Sb., 1988, **135**, 4, 472–496; English transl.: Math. USSR-Sb., 1989, **63**, 457–482.
- [5] E. Bese, *On the spannedness and very ampleness of certain line bundles on the blow-ups of $\mathbb{P}_{\mathbb{C}}^2$ and $\mathbb{F}_r$* , Math. Ann. **262** (1983), 225–238.
- [6] G. Brown, A. Corti, F. Zucconi *Birational geometry of 3-fold Mori fibre spaces*, Proceedings of the Fano Conference (Torino, Italy, 29 Sept.–5 Oct. 2002), A. Conte, A. Collino and M. Marchisio Eds., Torino, 2004, pp. 235–275.
- [7] I. Cheltsov, *Non-rational nodal quartic threefolds*, Pacific J. of Math., **226** (2006), 1, 65–82.
- [8] I. Cheltsov, *Points in projective spaces and applications*, arXiv:math.AG/0511578 (2006).
- [9] I. Cheltsov, J. Park, *Sextic double solids*, arXiv:math.AG/0404452 (2004).
- [10] C. Ciliberto, V. di Gennaro, *Factoriality of certain hypersurfaces of $\mathbb{P}^3$ with ordinary double points* Encyclopaedia of Mathematical Sciences **132** Springer-Verlag, Berlin, (2004), 1–9.
- [11] S. Cynk, *Defect of a nodal hypersurface*, Manuscripta Math. **104** (2001), 325–331.
- [12] D. Eisenbud, J.-H. Koh, *Remarks on points in a projective space*, Commutative algebra, Berkeley, CA (1987), MSRI Publications **15**, Springer, New York, 157–172.
- [13] K. Matsuki. *Introduction to the Mori program*. Universitext, Springer, 2002.

- [14] M. Mella, *Birational geometry of quartic 3-folds II: the importance of being $\mathbb{Q}$ -factorial*, Math. Ann. **330** (2004), 107–126.